

# BAPSoL™

PLANT GROWTH REGULATOR

ACTIVE CONSTITUENT: 20 g/L 6-BENZYLADENINE

**For Reducing Fruit Set, Increasing Fruit Size and Improving Firmness of Harvested Apples as Specified in the Directions For Use Table.**

## GENERAL INSTRUCTIONS

BAPSoL is a secondary thinner designed for use on trees treated with a primary thinner (ethephon or NAA) at or about flowering. Best results from BAPSoL will be obtained when foliage is evenly and thoroughly covered with the spray. Timing and temperature at time of application is critical. BAPSoL does not increase the incidence of russet when applied alone at label rates. An increase in vegetative growth may be observed following application of BAPSoL.

## MIXING

BAPSoL mixes easily with water. Add the required amount to a clean spray tank about half-filled with water. Agitate while bringing the total volume of water to the required level. Use the spray solution within 24 hours after mixing. Addition of a non-ionic wetting agent may improve coverage.

## APPLICATION

Apply in sufficient water to achieve thorough and even coverage of leaves. Coverage of lower leaf surfaces is important. Apply when king fruitlets are between 7 to 10 mm in diameter. This normally occurs 10 to 25 days after full bloom. Spray on a day when the daytime temperature is increasing to a predicted daily maximum of greater than 15 °C.

## COMPATIBILITY

It is recommended that BAPSoL not be mixed with other growth regulants, pesticides or fertilisers.

## PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

**DO NOT** apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops or pastures.

## PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

**DO NOT** contaminate streams, rivers or waterways with the chemical or used container.

## STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN.

Store in the closed, original container, in a well-ventilated area, as cool as possible.

**DO NOT** store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray

tank. **DO NOT** dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

## SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Avoid contact with eyes and skin.

When opening the container and preparing product for use wear cotton overalls buttoned to the neck and wrist and washable hat and elbow-length PVC gloves and face shield or goggles. When using the prepared spray wear cotton overalls buttoned to the neck and wrist, a washable hat and elbow-length PVC gloves. If product in eyes, wash it out immediately with water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

## FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

## SDS

Additional information is listed in the Material Safety Data Sheet (SDS).

## IMPORTANT NOTICE

These goods are to be used only for the purpose and as specified on the label, and are not suitable for any other purpose. To the fullest extent permitted by law, we do not accept or bear any liability on any basis for any loss, damage, cost or expense, arising in any way, directly or indirectly, in connection with the goods.

™ Trademark of Agrinova NZ Ltd

APVMA Approval No.: 58956/5/0904

58956/20/0904

In a Transport Emergency  
**Dial 000**  
Police or Fire Brigade

**SPECIALIST ADVICE  
IN EMERGENCY ONLY  
1800 024 973  
ALL HOURS - AUSTRALIA WIDE**

## DIRECTIONS FOR USE

### RESTRAINTS

**DO NOT** apply when maximum daily temperature is likely to be less than 15°C.

**DO NOT** apply later than 30 days after full bloom.

CROP	STATE	RATE	CRITICAL COMMENTS
<b>Apple varieties:</b> Fuji, Gala, Golden Delicious, Pink Lady, Red Delicious, Sundowner	All States	9 L/ha in a minimum of 1,200 L/ha	Use as a secondary thinner after application of a primary thinner (ethephon or NAA) at bloom. Increase water volume as canopy density increases. Apply when king fruitlets are between 7 to 10 mm in diameter. This normally occurs 10 to 25 days after full bloom. Spray on a day when the daytime temperature is increasing to a predicted daily maximum of greater than 15°C.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL  
UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

### WITHHOLDING PERIOD

**NOT REQUIRED** WHEN USED AS DIRECTED.